

Storm Prediction Center



Site Map

News Organization

Search for:

• SPC NCEP All NOAA Go

Local forecast by "City, St" or "ZIP"

City, St

Go

SPC on Facebook



@NWSSPC

NCEP Quarterly Newsletter

Home (Classic) **SPC Products**

All SPC Forecasts Current Watches Meso. Discussions **Conv. Outlooks Tstm. Outlooks Fire Wx Outlooks** RSS Feeds E-Mail Alerts Weather Information **Storm Reports**

Storm Reports Dev. NWS Hazards Map Watch/Warning Map National RADAR **Product Archive NOAA** Weather Radio

Research


Non-op. Products **Forecast Tools** Svr. Tstm. Events **SPC Publications** SPC-NSSL HWT

Education & Outreach

About the SPC SPC FAQ **About Tornadoes About Derechos Video Lecture Series WCM Page** Enh. Fujita Page **Our History Public Tours**

Misc. Staff **Contact Us SPC Feedback**

Mesoscale Discussion 1185



SPC MCD #1185

Mesoscale Discussion 1185 NWS Storm Prediction Center Norman OK 0226 PM CDT Thu Jun 29 2017

Areas affected...Central into northeast NE...Southeast SD...Northwest IA

Concerning...Severe potential...Watch likely

Valid 291926Z - 292100Z

Probability of Watch Issuance...80 percent

SUMMARY...At least isolated severe thunderstorm development appears likely later this afternoon. Large hail and damaging wind will be the primary threat. Watch issuance is likely for portions of the area.

DISCUSSION...Tomwering cumulus has developed along a cold front across portions of southwest into north-central NE. With time, scattered thunderstorm development is expected along the front, and



also in the vicinity of an outflow boundary running from northwest IA into northeast NE/southeast SD, as a midlevel vort max over western SD moves eastward through the afternoon. While the 18Z OAX sounding still exhibits the effects of the nocturnal MCS across KS/MO, more substantial recovery has likely occurred farther west, with strong heating and steep midlevel lapse rates supporting moderate MLCAPE of 1500-2500 J/kg, while westerly midlevel flow is sufficient to support effective shear of 35-45 kt.

Initial supercell development is possible both along the front and the outflow boundary. Isolated instances of very large hail in excess of 2 inches in diameter will be possible, along with isolated damaging wind gusts. A tornado or two also will be possible with any supercells that interact with the outflow boundary across northeast NE/northwest IA. With time, some upscale growth is possible with an increasing threat of damaging wind. Watch issuance is likely later this afternoon to cover these threats.

..Dean/Guyer.. 06/29/2017

...Please see www.spc.noaa.gov for graphic product...

ATTN...WFO...FSD...OAX...ABR...GID...LBF...GLD...

LAT...LON 40610065 41010039 41400002 42069947 42649927 43139928 43739946 43579812 43109682 42879613 42649602 42189620 41729679 41289760 40919839 40499924 40219960 40200005 40310045 40610065

Top/All Mesoscale Discussions/Forecast Products/Home

Weather Topics:

Watches, Mesoscale Discussions, Outlooks, Fire Weather, All Products, Contact Us

NOAA / National Weather Service National Centers for Environmental Prediction Storm Prediction Center 120 David L. Boren Blvd. Norman, OK 73072 U.S.A. spc.feedback@noaa.gov Page last modified: June 29, 2017

Disclaimer
Information Quality
Help
Glossary

Privacy Policy
Freedom of Information Act (FOIA)
About Us
Career Opportunities