



upslope flow. Effective bulk shear of 40-55 knots and forecast MLCAPE of 1500-2000 J/kg should result in a conducive environment for strong to severe thunderstorms. While low level flow remains relatively weak, easterly surface winds with stronger southwest flow aloft should provide enough veering in the profile to support rotating updrafts.

Latest HRRR guidance shows a few discrete supercells forming along and east of the Front Range and continuing eastward through the afternoon. A multi-cell cluster should form in southeast Colorado separate from the Front Range convection in the vicinity of a surface low and front. These could merge later and form a downstream MCS into the evening. Large hail, damaging winds, and a few tornadoes are all possible if not likely especially with the strongest storms.

..Nauslar/Peters/Weiss.. 06/19/2018

... Please see www.spc.noaa.gov for graphic product...

ATTN...WFO...LBF...GLD...AMA...PUB...BOU...CYS...

LAT...LON 41020550 40980514 40980364 41000228 40830211 39540215 37970211 37090215 36990267 37080374 37040439 37610479 38220503 38980501 39520511 40220525 40590536 41020550

Top/All Mesoscale Discussions/Forecast Products/Home

## Weather Topics:

Watches, Mesoscale Discussions, Outlooks, Fire Weather, All Products, Contact Us

NOAA / National Weather Service National Centers for Environmental Prediction Storm Prediction Center 120 David L. Boren Blvd. Norman, OK 73072 U.S.A. spc.feedback@noaa.gov Page last modified: June 19, 2018 Disclaimer Information Quality Help Glossary Privacy Policy Freedom of Information Act (FOIA) About Us Career Opportunities