

# Storm Prediction Center



Map

**News Organization** 

Search for:

• SPC NCEP All NOAA Go

## Local forecast by "City, St" or "ZIP"

City, St

Go





@NWSSPC

**NCEP Quarterly Newsletter** 

## Home (Classic) **SPC Products**

**All SPC Forecasts Current Watches** Meso. Discussions Conv. Outlooks **Tstm. Outlooks Fire Wx Outlooks** RSS Feeds **E-Mail Alerts Weather Information Storm Reports Storm Reports Dev. NWS Hazards Map National RADAR Product Archive NOAA** Weather Radio

#### Research

Non-op. Products **Forecast Tools** Svr. Tstm. Events **SPC Publications** SPC-NSSL HWT


## **Education & Outreach About the SPC**

**SPC FAQ About Tornadoes** About Derechos Video Lecture Series **WCM Page** Enh. Fujita Page **Our History Public Tours** 

Misc. Staff

**Contact Us SPC Feedback** 

## **Mesoscale Discussion 572**



SPC MCD #0572

Mesoscale Discussion 0572 NWS Storm Prediction Center Norman OK 0249 PM CDT Tue May 07 2019

Areas affected...Parts of northeastern New Mexico and southeastern Colorado

Concerning...Severe potential...Watch possible

Valid 071949Z - 072145Z

Probability of Watch Issuance...40 percent

SUMMARY...Scattered strong to severe storms, including isolated supercells, appear likely to develop off the Sangre de Cristo Mountains, across and south of the Raton Mesa area through 3-5 PM MDT. It is not yet certain that a watch will be required, but trends are being monitored for this possibility.

DISCUSSION...Downstream of the closed mid-level low now turning east-northeastward into the southern Rockies, orographic and



large-scale forcing for ascent are contributing to the initiation of thunderstorms near the Sangre de Cristo Mountains. At the same time, insolation within post-frontal easterly low-level to the lee of the mountains is contributing to steepening low-level lapse rates and weak to modest boundary-layer destabilization. As inhibition for boundary layer parcels continues to erode, storms will propagate east of the mountain through late afternoon, aided by 30+ kt southwesterly ambient mean flow. With pronounced veering of lower through mid-tropospheric wind fields contributing to strong deep layer shear, a few supercells are possible, accompanied primarily by the risk for severe hail and localized strong surface gusts.

..Kerr/Guyer.. 05/07/2019

...Please see www.spc.noaa.gov for graphic product...

ATTN...WFO...PUB...ABQ...

LAT...LON 37250434 37820433 37170363 36900319 35200320 35140433 35930460 36900450 37250434

### Top/All Mesoscale Discussions/Forecast Products/Home

### Weather Topics:

Watches, Mesoscale Discussions, Outlooks, Fire Weather, All Products, Contact Us

NOAA / National Weather Service
National Centers for Environmental Prediction
Storm Prediction Center
120 David L. Boren Blvd.
Norman, OK 73072 U.S.A.
spc.feedback@noaa.gov
Page last modified: May 07, 2019

Disclaimer
Information Quality
Help
Glossary

Privacy Policy
Freedom of Information Act (FOIA)
About Us
Career Opportunities