

## **Storm Prediction Center**



Local forecast by
"City, St" or "ZIP"
City, St

Site Map

Find us on Facebook

**@NWSSPC** 

NCEP Quarterly Newsletter

Home (Classic) **SPC Products All SPC Forecasts Current Watches Meso. Discussions Conv. Outlooks Tstm. Outlooks Fire Wx Outlooks** NSS Feeds **E-Mail Alerts** Weather Information **Storm Reports Storm Reports Dev. NWS Hazards Map National RADAR Product Archive** NOAA Weather Radio

Research Non-op. Products **Forecast Tools Svr. Tstm. Events SPC Publications SPC-NSSL HWT** Education & Outreach **About the SPC SPC FAQ About Tornadoes About Derechos Video Lecture Series WCM Page** Enh. Fujita Page **Our History Public Tours** 



**SPC Feedback** 


Misc.

Staff Contact Us

**Mesoscale Discussion 1327** 

News

**Organization** 



Mesoscale Discussion 1327 NWS Storm Prediction Center Norman OK 1201 AM CDT Thu Jun 29 2023

Areas affected...Northeast CO into southwest NE/northwest KS

Concerning...Severe Thunderstorm Watch 410...

Valid 290501Z - 290630Z

The severe weather threat for Severe Thunderstorm Watch 410 continues.

SUMMARY...A threat for very large hail may continue into the early morning hours with ongoing supercells. There is some potential for the severe-gust potential to increase with time.

DISCUSSION...A large and intense supercell is ongoing at 0445 UTC across extreme northeast CO, with other more recent development noted near/north of the trailing outflow into a larger part of northeast CO. In the short term, moderate to strong buoyancy and favorable deep-layer shear will continue to support supercell potential, with an attendant threat of very large hail. A short-term tornado threat also cannot be ruled out with the lead supercell or with any of the trailing supercells that can remain rooted near the surface and not be undercut by outflow.

The lead supercell has shown an increase in forward speed over the last hour, and there is some potential for this cell to evolve into a forward-propagating cluster that could persist for much of the night. The evolution remains uncertain, however, due to the likely somewhat elevated nature of any overnight cluster, and the potentially negative impact of very warm temperatures at the base of the EML. If an organized cluster can persist and propagate eastward, a swath of severe gusts (in addition to hail) will be possible overnight, and downstream watch issuance may eventually be needed.

- ..Dean/Thompson.. 06/29/2023
- ...Please see www.spc.noaa.gov for graphic product...

ATTN...WFO...GID...LBF...GLD...BOU...

LAT...LON 40730440 40950188 40940173 40950051 40899968 40619917 40239894 39889906 39649957 39610062 39590137 39630273 39660344 39800410 40000435 40080495 40730440

Top/All Mesoscale Discussions/Forecast Products/Home

Weather Topics:
Watches, Mesoscale Discussions, Outlooks, Fire Weather, All Products, Contact Us

NOAA / National Weather Service National Centers for Environmental Prediction Storm Prediction Center 120 David L. Boren Blvd. Norman, OK 73072 U.S.A. spc.feedback@noaa.gov

Page last modified: June 29, 2023

Disclaimer Information Quality Help Glossary Privacy Policy Freedom of Information Act (FOIA) About Us Career Opportunities

• SPC NCEP All NOAA Go

Search for: